

STRONGLY RIGHT ALTERNATIVE RINGS AND BOL LOOPS

MICHAEL K. KINYON, KENNETH KUNEN*, AND J. D. PHILLIPS

ABSTRACT. We partially answer two questions of Goodaire by showing that in a finite, strongly right alternative ring, the set of units (if the ring is with unity) is a Bol loop under ring multiplication, and the set of quasiregular elements is a Bol loop under “circle” multiplication.

A *magma* $(L, \cdot)$ is a set L with a binary operation $\cdot : L \times L \rightarrow L$. A magma $(L, \cdot)$ is a *loop* if there is a neutral element $1 \in L$ satisfying $1x = x1 = x$, and if all right translations $R(a) : x \mapsto xa$ and all left translations $L(a) : x \mapsto ax$ are bijections. If a magma L satisfies the *right Bol identity* $(xy \cdot z)y = x(yz \cdot y)$, then we will say that L is a *Bol magma*. The Bol identity can be equivalently expressed in terms of right translations: $R(y)R(z)R(y) = R(yz \cdot y)$. A *Moufang loop* is a Bol loop which also satisfies the *flexible law* $x \cdot yx = xy \cdot x$. For background in loops, the reader is referred to the standard references [1] and [9], and further details on Bol loops can be found in [10].

A (nonassociative) ring $(R, +, \cdot)$ is *right alternative* if it satisfies the *right alternative law* $xy \cdot y = xy^2$. A right alternative ring R is *strongly right alternative* if $(R, \cdot)$ is a Bol magma. A right alternative ring of characteristic not 2 is strongly right alternative ([11], §16.1), but this is not true in general [3] [7]. An *alternative* ring is a right alternative ring which also satisfies the *left alternative law* $x \cdot xy = x^2y$. An alternative ring is strongly right alternative ([11], §2.3), but there are strongly right alternative rings which are not alternative.

In a ring R with unity, a *unit* is an element with a (not necessarily unique) two-sided inverse. Let $\mathcal{U}(R)$ denote the set of all units in R . It is well-known that if R is an associative ring with unity, then $\mathcal{U}(R)$ is a group. More generally, if R is an alternative ring with unity, then $\mathcal{U}(R)$ is a Moufang loop ([5], §11.5.3). In a strongly right alternative ring with unity, it is known that if $\mathcal{U}(R)$ is closed under multiplication, then $\mathcal{U}(R)$ is a Bol loop ([4], Lem. 3.1, p. 356). Goodaire posed the following problem ([4], p. 356):

Problem 1. *In a strongly right alternative ring R with unity, is $\mathcal{U}(R)$ a Bol loop?*

In a ring R , define a binary operation $\circ : R \times R \rightarrow R$ by $x \circ y = x + y + x \cdot y$ for $x, y \in R$. An element x is said to be *quasiregular* with *quasi-inverse* x' if $x \circ x' = x' \circ x = 0$. Let $\mathcal{Q}(R)$ denote the set of all quasiregular elements of R . If R is

1991 *Mathematics Subject Classification.* 17D05, 20N05.

Key words and phrases. right alternative ring, quasiregular element, Bol loop.

*Partially supported by NSF Grant DMS-0456653.

associative, then $(R, \circ)$ is associative, and so $(\mathcal{Q}(R), \circ)$ is a group. More generally, if R is alternative, then $(\mathcal{Q}(R), \circ)$ is a Moufang loop [2]. For a right alternative ring R , the magma $(R, \circ)$ is right alternative, that is, $(x \circ y) \circ y = x \circ (y \circ y)$, and if R is strongly right alternative, then $(R, \circ)$ is a Bol magma with neutral element 0 ([3], Lem. 3.1). Thus we have the following problem [3].

Problem 2. *In a strongly right alternative ring R , is $(\mathcal{Q}(R), \circ)$ a Bol loop?*

In this paper, we offer partial affirmative answers to Problems 1 and 2. Our main result, however, is not actually about right alternative rings. In what follows, let $(L, \cdot)$ denote a Bol magma with neutral element $1 \in L$, and let $\mathcal{J}(L)$ denote the subset of all elements of L with two-sided inverses, that is, for each $a \in \mathcal{J}(L)$, there exists $a^{-1} \in L$ such that $aa^{-1} = a^{-1}a = 1$. (The uniqueness of a^{-1} will be addressed in Lemma 4 below.)

Theorem 1. *If L is a finite Bol magma with a neutral element, then $(\mathcal{J}(L), \cdot)$ is a Bol loop.*

Implicit in the statement of Theorem 1 is the assertion that $\mathcal{J}(L)$ is closed under multiplication. In fact by the following lemma, the closure is all that needs to be verified.

Lemma 1. *If L is a Bol magma with a neutral element, and if $\mathcal{J}(L)$ is closed under multiplication, then $(\mathcal{J}(L), \cdot)$ is a loop.*

A proof of Lemma 1 can be extracted from the proof of ([4], Lem. 3.1, p. 356). We will give an independent proof for completeness.

As an immediate consequence of Theorem 1, we obtain our partial affirmative answers to Problems 1 and 2.

Corollary 1. *Let $(R, +, \cdot)$ be a finite, strongly right alternative ring.*

- (1) *If R is with unity, then $(\mathcal{U}(R), \cdot)$ is a Bol loop.*
- (2) *$(\mathcal{Q}(R), \circ)$ is a Bol loop.*

We will also use Lemma 1 to obtain an auxiliary result.

Theorem 2. *If L is a flexible Bol magma with a neutral element, then $(\mathcal{J}(L), \cdot)$ is a Moufang loop.*

As a consequence, we obtain a new proof of the following; see §11.5.3 of [5] and [2].

Corollary 2. *Let $(R, +, \cdot)$ be an alternative ring.*

- (1) *If R is with unity, then $(\mathcal{U}(R), \cdot)$ is a Moufang loop.*
- (2) *$(\mathcal{Q}(R), \circ)$ is a Moufang loop.*

Our investigations were aided by the automated deduction tool OTTER developed by McCune [8].

Recall the definition of powers in a magma: $a^0 := 1$, $a^n := a^{n-1}a$ for all positive integers n .

Lemma 2. *For all nonnegative integers n , and all $a \in L$, $R(a)^n = R(a^n)$.*

Proof. The proof of this result for Bol loops carries through to the present setting without change; see [10] or ([9], IV.6.5). $\square$

Lemma 3. *If $a, b, c \in L$ satisfy $ab = ca = 1$, then $b = c$.*

Proof. For all x , the right Bol identity gives $c(ax \cdot a) = xa$. Taking $x = b^2$ and using Lemma 2, we get $c \cdot ba = b^2a$. Taking $x = b$, we obtain $ba = 1$. Thus $c = b^2a = (ba \cdot b^2)a = b(ab^2 \cdot a) = b \cdot ba = b$, using Lemma 2 and the right Bol identity again. The rest is clear. $\square$

Lemma 4. *For each $a \in \mathcal{J}(L)$,*

1. *There is a unique two-sided inverse $a^{-1} \in \mathcal{J}(L)$,*
2. *$R(a)$ is a bijection of L and $R(a)^{-1} = R(a^{-1})$,*
3. *$L(a)$ is a bijection of L and $L(a)^{-1} = R(a)L(a^{-1})R(a^{-1})$.*

Proof. (1) is immediate from Lemma 3. For (2), we compute

$$R(a)R(a^{-1})R(a) = R(aa^{-1} \cdot a) = R(a) \quad (1)$$

by the right Bol identity. Next, set $c := (a^{-1})^2$ and compute

$$R(a)R(c)R(a) = R(ac \cdot a) = I \quad (2)$$

since $ac = aa^{-1} \cdot a^{-1}$ by Lemma 2. Thus

$$R(a^{-1})R(a) = R(a)R(c)R(a)R(a^{-1})R(a) = R(a)R(c)R(a) = I$$

using (2), (1), and (2) again. A similar argument gives $R(a)R(a^{-1}) = I$.

For (3): The right Bol identity can be written as $L(xy)R(y) = L(y)R(y)L(x)$. Taking $x = a^{-1}$, $y = a$, and using $R(a)^{-1} = R(a^{-1})$, we have

$$L(a)[R(a)L(a^{-1})R(a^{-1})] = I.$$

On the other hand, taking $x = a$ and $y = a^{-1}$, then $[R(a)L(a^{-1})R(a^{-1})]L(a) = I$. This completes the proof. $\square$

The proof of part (2) of Lemma 4 is an adaptation to this setting of the proof of Theorem 3.10(3) of [6] (see also Remark 5, p. 51).

We now prove our main lemma.

Proof of Lemma 1. If $\mathcal{J}(L)$ is closed under multiplication, then Lemma 4 implies that for each $a \in \mathcal{J}(L)$, the restrictions of $R(a)$ and $L(a)$ to $\mathcal{J}(L)$ are bijections. Therefore $\mathcal{J}(L)$ is a loop. $\square$

Lemma 5. *For each $a, b \in \mathcal{J}(L)$, $L(ab)$ is bijective, and*

$$L(ab)^{-1} = R(b)R(a)L(b^{-1}a^{-1})R(a^{-1})R(b^{-1}).$$

In particular, ab has a unique right inverse, that is, there exists a unique $c \in L$ such that $ab \cdot c = 1$.

Proof. By the Bol identity, $L(ab)R(b) = L(b)R(b)L(a)$, and so by Lemma 4 and the Bol identity again,

$$\begin{aligned} L(ab)^{-1} &= R(b)L(a)^{-1}R(b^{-1})L(b)^{-1} \\ &= R(b)R(a)L(a^{-1})R(a^{-1})R(b^{-1})R(b)L(b^{-1})R(b^{-1}) \\ &= R(b)R(a)L(a^{-1})R(a^{-1})L(b^{-1})R(b^{-1}) \\ &= R(b)R(a)L(b^{-1}a^{-1})R(a^{-1})R(b^{-1}) \end{aligned}$$

The right inverse of ab is $c := 1L(ab)^{-1} = (b^{-1}a^{-1} \cdot ba)a^{-1} \cdot b^{-1}$. $\square$

Lemma 6. *Suppose $a, b \in L$ satisfy $ab = 1$, and suppose there exist distinct positive integers m, n such that $b^m = b^n$. Then $ba = 1$.*

Proof. By Lemma 2, $R(b)^m = R(b)^n$. Applying this to a , we obtain $R(b)^{m-1} = R(b)^{n-1}$. Hence there exists a positive integer k such that $R(b)^k = I$, i.e., $b^k = 1$. Thus b has a right inverse which, by Lemma 3, must be a . $\square$

Finally, we turn to the proof of our main result.

Proof of Theorem 1. Fix $a, b \in \mathcal{J}(L)$. By Lemma 5, ab has a right inverse c . Since L is finite, there exist distinct positive integers m, n such that $c^m = c^n$. By Lemma 6, $ab \in \mathcal{J}(L)$. By Lemma 1, $\mathcal{J}(L)$ is a loop. $\square$

As the proofs of Lemma 6 and Theorem 1 indicate, the finiteness of L can obviously be relaxed to a finite torsion assumption: for each $a \in L$, there exist distinct positive integers m, n (depending on a) such that $a^m = a^n$.

Next we turn to our auxiliary result.

Proof of Theorem 2. Fix $a, b \in \mathcal{J}(L)$, and let $c \in \mathcal{J}(L)$ be the right inverse of ab given by Lemma 5. Then $ab \cdot (c \cdot ab) = (ab \cdot c) \cdot ab = ab = ab \cdot 1$. By Lemma 5, we may cancel ab on the left to obtain $c \cdot ab = 1$ as claimed. Thus $ab \in \mathcal{J}(L)$, and so by Lemma 1, $\mathcal{J}(L)$ is a loop. $\square$

Proof of Corollary 2. Let $(R, +, \cdot)$ be an alternative ring. If R is with unity, then R is strongly right alternative and $(R, \cdot)$ is flexible ([11], §2.3). Thus $(\mathcal{U}(R), \cdot)$ is a flexible Bol magma with a neutral element, and so by Theorem 2, $(\mathcal{U}(R), \cdot)$ is a Moufang loop. Also, $(R, \circ)$ is a flexible Bol magma with neutral element 0 ([2], [3], Lem. 3.1), and so Theorem 2 implies that $(\mathcal{Q}(R), \circ)$ is a Moufang loop. $\square$

By applying Lemma 1, one may show that $\mathcal{J}(L)$ is a loop for various other special classes of Bol magmas, but lacking examples or motivation, we omit these. In general, we suspect the following to be the case.

Conjecture. *There exists a Bol magma L with a neutral element such that $\mathcal{J}(L)$ is not closed under multiplication.*

Returning to strongly right alternative rings, it is possible that the ring structure allows further weakening of the torsion assumptions on $(\mathcal{U}(R), \cdot)$ or $(\mathcal{Q}(R), \circ)$

in Corollary 1. In any case, we have not been able to find a (necessarily infinite) strongly right alternative ring R such that either $(\mathcal{U}(R), \cdot)$ or $(\mathcal{Q}(R), \circ)$ is not closed under multiplication. Thus Problems 1 and 2 remain open.

REFERENCES

- [1] R. H. Bruck, *A Survey of Binary Systems*, Springer-Verlag, 1971.
- [2] E. G. Goodaire, Circle loops of radical alternative rings, *Algebras Groups Geom.* **4** (1987), 461–474.
- [3] E. G. Goodaire, Nilpotent right alternative rings and Bol circle loops, *Comm. Algebra* **28** (2000), 2445–2459.
- [4] E. G. Goodaire, Units in right alternative loop rings, *Publ. Math. Debrecen* **59** (2001), 353–362.
- [5] E. G. Goodaire, E. Jespers, and C. Polcino Milies, *Alternative Loop Rings*, vol. **184**, North-Holland Math. Studies, Elsevier, 1996.
- [6] H. Kiechle, *Theory of K-Loops*, Lecture Notes in Math. **1779**, Springer-Verlag, 2002.
- [7] K. Kunen, Alternative loop rings, *Comm. Algebra* **26** (1998), 557–564.
- [8] W. W. McCune, *OTTER 3.3 Reference Manual and Guide*, Technical Memorandum ANL/MCS-TM-263, Argonne National Laboratory, 2003; or see:
<http://www.mcs.anl.gov/AR/otter/>
- [9] H. O. Pflugfelder, *Quasigroups and Loops: Introduction*, Sigma Series in Pure Math. **8**, Heldermann Verlag, Berlin, 1990.
- [10] D. A. Robinson, Bol loops, *Trans. Amer. Math. Soc.* **123** (1966), 341–354.
- [11] K. A. Zhevlakov, A. M. Slin'ko, I. P. Shestakov, and A. I. Shirshov, *Rings that are Nearly Associative*, Academic Press, New York, 1982, trans. H. F. Smith.

DEPARTMENT OF MATHEMATICAL SCIENCES, INDIANA UNIVERSITY SOUTH BEND, SOUTH BEND, IN 46634 USA

E-mail address: mkinyon@iusb.edu

URL: <http://www.iusb.edu/~mkinyon>

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF WISCONSIN, MADISON, WI 57306 USA

E-mail address: kunen@math.wisc.edu

URL: <http://www.math.wisc.edu/~kunen>

DEPARTMENT OF MATHEMATICS & COMPUTER SCIENCE, WABASH COLLEGE, CRAWFORDSVILLE, IN 47933 USA

E-mail address: phillipj@wabash.edu

URL: <http://www.wabash.edu/depart/math/faculty.html#Phillips>